

Chromatic number and spectral radius

Vladimir Nikiforov

Department of Mathematical Sciences, University of Memphis,
Memphis TN 38152, USA; e-mail: vnikifrv@memphis.edu

February 2, 2008

Abstract

Write $\mu(A) = \mu_1(A) \geq \dots \geq \mu_{\min}(A)$ for the eigenvalues of a Hermitian matrix A . Our main result is: let A be a Hermitian matrix partitioned into $r \times r$ blocks so that all diagonal blocks are zero. Then for every real diagonal matrix B of the same size as A ,

$$\mu(B - A) \geq \mu\left(B + \frac{1}{r-1}A\right).$$

Let G be a nonempty graph, $\chi(G)$ be its chromatic number, A be its adjacency matrix, and L be its Laplacian. The above inequality implies the well-known result of A.J. Hoffman

$$\chi(G) \geq 1 + \frac{\mu(A)}{-\mu_{\min}(A)},$$

and also,

$$\chi(G) \geq 1 + \frac{\mu(A)}{\mu(L) - \mu(A)}.$$

Equality holds in the latter inequality if and only if every two color classes of G induce a $|\mu_{\min}(A)|$ -regular subgraph.

Keywords: *graph Laplacian; largest eigenvalue; least eigenvalue; k -partite graph; chromatic number.*

AMS classification: 05C50.

Write $\mu(A) = \mu_1(A) \geq \dots \geq \mu_{\min}(A)$ for the eigenvalues of a Hermitian matrix A . Given a graph G , let $\chi(G)$ be its chromatic number, $A(G)$ be its adjacency matrix, and $D(G)$ be the diagonal matrix of its degree sequence; set $L(G) = D(G) - A(G)$.

Letting G be a nonempty graph with $L(G) = L$ and $A(G) = A$, we prove that

$$\chi(G) \geq 1 + \frac{\mu(A)}{\mu(L) - \mu(A)}, \tag{1}$$

complementing the well-known inequality of A.J. Hoffman [1]

$$\chi(G) \geq 1 + \frac{\mu(A)}{-\mu_{\min}(A)}. \quad (2)$$

Equality holds in (1) if and only if every two color classes of G induce a $|\mu_{\min}(A)|$ -regular subgraph.

We deduce inequalities (1) and (2) from a theorem of its own interest.

Theorem 1 *Let A be a Hermitian matrix partitioned into $r \times r$ blocks so that all diagonal blocks are zero. Then for every real diagonal matrix B of the same size as A ,*

$$\mu(B - A) \geq \mu\left(B + \frac{1}{r-1}A\right). \quad (3)$$

Proof of Theorem 1 Write n for the size of A , let $[n] = \cup_{i=1}^r N_i$ be the partition of its index set, and let $b_1, \dots, b_n$ be the diagonal entries of B . Set $L = B - A$, $K = (r-1)B + A$, and select a unit eigenvector $\mathbf{x} = (x_1, \dots, x_n)$ to $\mu(K)$. Our proof strategy is simple: using $\mathbf{x}$, we define specific n -vectors $\mathbf{y}_1, \dots, \mathbf{y}_r$ and show that

$$r(r-1)\mu(L) \geq \mu(L) \sum_{i \in [r]} \|\mathbf{y}_i\|^2 \geq \sum_{i \in [r]} \langle L\mathbf{y}_i, \mathbf{y}_i \rangle \geq r \langle K\mathbf{x}, \mathbf{x} \rangle = r\mu(K).$$

For $i = 1, \dots, r$ define $\mathbf{y}_i = (y_{i1}, \dots, y_{in})$ as

$$y_{ij} = \begin{cases} -(r-1)x_j & \text{if } j \in N_i \\ x_j & \text{if } j \in [n] \setminus N_i. \end{cases}$$

The Rayleigh principle implies that

$$\mu(L) \sum_{i \in [r]} \|\mathbf{y}_i\|^2 \geq \sum_{i \in [r]} \mu(L) \|\mathbf{y}_i\|^2 \geq \sum_{i \in [r]} \langle L\mathbf{y}_i, \mathbf{y}_i \rangle. \quad (4)$$

Noting that

$$\|\mathbf{y}_i\|^2 = \sum_{j \in [n] \setminus N_i} |x_j|^2 + (r-1)^2 \sum_{j \in N_i} |x_j|^2 = 1 + r(r-2) \sum_{j \in N_i} |x_j|^2,$$

we obtain,

$$\sum_{i \in [r]} \|\mathbf{y}_i\|^2 = r + r(r-2) \sum_{i \in [r]} \sum_{j \in N_i} |x_j|^2 = r + r(r-2) = r(r-1). \quad (5)$$

On the other hand, we have

$$\langle L\mathbf{y}_i, \mathbf{y}_i \rangle = \sum_{j \in [n]} b_j |y_{ij}|^2 - \sum_{j,k \in [n]} a_{jk} y_{ik} \overline{y_{ij}}.$$

For every $i \in [n]$, we see that

$$\sum_{j \in [n]} b_j |y_{ij}|^2 = \sum_{j \in [n]} b_j |x_j|^2 + r(r-2) \sum_{j \in N_i} b_j |x_j|^2,$$

and, likewise,

$$\sum_{j,k \in [n]} a_{jk} y_{ik} \overline{y_{ij}} = \sum_{j,k \in [n]} a_{jk} x_k \overline{x_j} - r \sum_{j \in N_i, k \in [n]} a_{jk} x_k \overline{x_j} - r \sum_{k \in N_i, j \in [n]} a_{jk} x_k \overline{x_j}.$$

Summing these for all $i \in [r]$, we find that

$$\begin{aligned} \sum_{i \in [r]} \langle L\mathbf{y}_i, \mathbf{y}_i \rangle &= \sum_{i \in [r], j \in [n]} b_j |x_j|^2 + r(r-2) \sum_{i \in [r], j \in N_i} b_j |x_j|^2 \\ &\quad - r \sum_{j,k \in [n]} a_{jk} x_k \overline{x_j} + r \sum_{i \in [r]} \left(\sum_{j \in N_i, k \in [n]} a_{jk} x_k \overline{x_j} + r \sum_{k \in N_i, j \in [n]} a_{jk} x_k \overline{x_j} \right) \\ &= r(r-1) \sum_{j \in [n]} b_j |x_j|^2 - r \sum_{j,k \in [n]} a_{jk} x_k \overline{x_j} + 2r \sum_{j,k \in [n]} a_{jk} x_k \overline{x_j} \\ &= r(r-1) \sum_{j \in [n]} b_j |x_j|^2 + r \sum_{j,k \in [n]} a_{jk} x_k \overline{x_j} = r \langle K\mathbf{x}, \mathbf{x} \rangle = r\mu(K). \end{aligned}$$

Hence, in view of (4) and (5), we obtain $(r-1)\mu(B-A) \geq \mu(K)$, completing the proof. $\square$

Lemma 2 *Let A be an irreducible nonnegative symmetric matrix and R be the diagonal matrix of its rowsums. Then*

$$\mu\left(R + \frac{1}{r-1}A\right) \geq \frac{r}{r-1}\mu(A) \quad (6)$$

with equality holding if and only if all rowsums of A are equal.

Proof Let $A = (a_{ij})$ and n be its size. Note first that for any vector $\mathbf{x} = (x_1, \dots, x_n)$

$$\langle (R-A)\mathbf{x}, \mathbf{x} \rangle = \sum_{1 \leq i < j \leq n} a_{ij} (x_i - x_j)^2 \geq 0.$$

Hence, $R-A$ is positive semidefinite; since A is irreducible, if $\langle (R-A)\mathbf{x}, \mathbf{x} \rangle = 0$, then all entries of $\mathbf{x}$ are equal.

Let $\mathbf{x} = (x_1, \dots, x_n)$ be an eigenvector to $\mu = \mu\left(D + \frac{1}{r-1}A\right)$. We have

$$\begin{aligned} \mu &= \left\langle \left(R + \frac{1}{r-1}A\right)\mathbf{x}, \mathbf{x} \right\rangle = \sum_{i=1}^n x_i^2 \sum_{j=1}^n a_{ij} + \frac{1}{r-1} \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j \\ &= \sum_{1 \leq i < j \leq n} a_{ij} (x_i - x_j)^2 + \frac{r}{r-1} \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j \\ &= \langle (R-A)\mathbf{x}, \mathbf{x} \rangle + \frac{r}{r-1} \sum_{i=1}^n \sum_{j=1}^n a_{ij} x_i x_j \geq \frac{r}{r-1} \mu(A), \end{aligned} \quad (7)$$

proving (6).

Let now equality holds in (6). Then equality holds in (7), and so $\langle (R - A)\mathbf{x}, \mathbf{x} \rangle = 0$ and $\mathbf{x}$ is an eigenvector of A to $\mu(A)$. Therefore $x_1 = \dots = x_n$ and the rowsums of A are equal.

If the rowsums of A are equal, the vector $\mathbf{j} = (1, \dots, 1)$ is an eigenvector of A to $\mu(A)$ and of R to $\mu(R)$; therefore $\mathbf{j}$ is an eigenvector of $R + \frac{1}{r-1}A$ to μ , and so equality holds in (6), completing the proof. $\square$

Proof of (1) and (2) Let G be a graph with chromatic number $\chi = r$. Coloring the vertices of G into r colors defines a partition of its adjacency matrix $A = A(G)$ with zero diagonal blocks. Letting B be the zero matrix, Theorem 1 implies inequality (2).

Letting now $B = D = D(G)$, Lemma 2 implies that

$$\mu\left(D + \frac{1}{r-1}A\right) \geq \frac{r}{r-1}\mu(A),$$

and inequality (1) follows.

The following argument for equality in (1) was kindly suggested by the referee. If equality holds in (1), by Lemma 2, G is regular; hence equality holds also in (2). Setting $\mu(G) = k$, $|\mu_{\min}(G)| = \tau$ and writing $\alpha(G)$ for the independence number of G , let us recall Hoffman's bound on $\alpha(G)$: for every k -regular graph G ,

$$\alpha(G) \leq \frac{n\tau}{k + \tau}. \quad (8)$$

On the other hand, we have

$$\alpha(G) \geq \frac{n}{\chi(G)} = \frac{n}{1 + k/\tau} = \frac{n\tau}{k + \tau}.$$

and thus, equality holds in (8). It is known (see, e.g., [2], Lemma 9.6.2) that this is only possible if $\chi(G) = n/\alpha(G)$ and every two color classes of G induce a τ -regular bipartite subgraph. $\square$

Concluding remarks

For the complete graph of order n without an edge, inequality (1) gives $\chi = n - 1$, while (2) gives only $\chi \geq n/2 + 2$. By contrast, for a sufficiently large wheel $W_{1,n}$, i.e., a vertex joined to all vertices of a cycle of length n , we see that (1) gives $\chi \geq 2$, while (2) gives $\chi \geq 3$.

A natural question is to determine when equality holds in (3). A particular answer, building upon [4], can be found in [5]: if G is a connected graph, then $\mu(D(G) - A(G)) = \mu(D(G) + A(G))$ if and only if G is bipartite.

Problem 3 *Determine when equality holds in (3).*

Finally, any lower bound on $\mu(A(G))$, together with (1), gives a lower bound on $\mu(L(G))$. This approach helps deduce some inequalities for bipartite graphs given in [3] and [6].

Acknowledgement. Thanks to Peter Rowlinson, Sebi Cioabă and Cecil Rousseau for useful suggestions. The author is most indebted to the referee for the exceptionally thorough, helpful and kind report.

References

- [1] A.J. Hoffman, On eigenvalues and colorings of graphs, *Graph Theory and its Applications*, Academic Press, New York (1970), pp. 79–91.
- [2] C. Godsil, G. Royle, *Algebraic Graph Theory*, Graduate Texts in Mathematics, 207, Springer-Verlag, New York, 2001, xx+439 pp.
- [3] Y. Hong, X.-D. Zhang, Sharp upper and lower bounds for largest eigenvalue of the Laplacian matrices of trees, *Discrete Math.* **296** (2005), 187-197.
- [4] R. Merris, Laplacian matrices of graphs: a survey, *Linear Algebra Appl.* **197–198** (1994), 143–176.
- [5] J.-L. Shu, Y. Hong, K. Wen-Ren, A sharp upper bound on the largest eigenvalue of the Laplacian matrix of a graph, *Linear Algebra Appl.* **347** (2002) 123–129.
- [6] A.M. Yu, M. Lu, F. Tian, On the spectral radius of graphs, *Linear Algebra Appl.* **387** (2004) 41–49.